

RATIONAL FUNCTIONS

Remember what we call a number like $\frac{2}{7}$? This is called a *rational* number because it is the *ratio* of two integers. In a like manner, a rational function is the ratio of two special functions called polynomials. Since a rational function is essentially a fraction, we will have to avoid dividing by zero, which means the domain may not be all real numbers.



□ POLYNOMIAL FUNCTIONS

Each of the following is a polynomial function:

$$y = 7 \quad \text{(a *linear* function -- it's a horizontal line)}$$

$$y = -3x + \sqrt{2} \quad \text{(a *linear* function -- it's a line with slope -3)}$$

$$y = 2x^2 - x + 9 \quad \text{(a *quadratic* function -- it's a parabola)}$$

$$f(x) = \sqrt[4]{2}x^3 - x^2 \quad \text{(a *cubic* function)}$$

$$P(x) = -\pi x^4 + 5x^2 + 8 \quad \text{(a *quartic* function)}$$

$$Q(x) = \frac{2}{3}x^5 + 1 \quad \text{(a *quintic* function)}$$

The key to any ***polynomial function*** is that all the exponents on the x come from \mathbb{W} , the set of whole numbers $\{0, 1, 2, 3, \dots\}$. The coefficients (the numbers in front of the variables), on the other hand, can come from anywhere in \mathbb{R} , the set of real numbers.

2

Consider the quartic polynomial

$$y = -2\pi x^4 + \frac{9}{10}x^3 - 17x^2 + \sqrt{2}.$$

First look at the exponents -- they're all whole numbers. Even the last term of the polynomial, $\sqrt{2}$, can be written as $\sqrt{2}x^0$, and so even the exponent on this last term is a whole number. Thus, all the exponents (4, 3, 2, and 0) come from \mathbb{W} , while all the coefficients (-2π , $\frac{9}{10}$, -17 , $\sqrt{2}$) come from \mathbb{R} . Considering the definition of polynomial, the given function is indeed a polynomial.

Each of the following is not a polynomial:

$$y = \frac{1}{x} \quad \left(\frac{1}{x} = x^{-1} \text{ and } -1 \notin \mathbb{W}\right)$$

$$y = \sqrt{x} \quad \left(\sqrt{x} = x^{1/2} \text{ and } \frac{1}{2} \notin \mathbb{W}\right)$$

$$f(x) = \frac{1}{\sqrt[3]{x}} \quad \left(\frac{1}{\sqrt[3]{x}} = x^{-1/3} \text{ and } -\frac{1}{3} \notin \mathbb{W}\right)$$

$$g(x) = |x - 1| \quad (\text{no absolute values allowed around the } x)$$

$$E(x) = 2^x \quad (\text{since } x \text{ is in the exponent, it can be any number})$$

$$T(x) = \sin x \quad (\text{it's on your calculator, but it's not a polynomial})$$

$$y = \log x \quad (\text{we'll learn this later -- it's not a polynomial})$$

$$x^2 + y^2 = 25 \quad (\text{it's a circle -- it's not a function of any kind})$$

Homework


1. Explain why $y = \pi x^5 - \sqrt{2}x^3 + \frac{1}{4}x - 17.5$ is a polynomial function.
2. Explain why $h(x) = \sqrt[3]{5}x^4 - \sqrt{2}x + \frac{1}{2}$ is not a polynomial function.

Intercepts come next. If $x = 0$, then $y = \frac{1}{0-2} = -\frac{1}{2}$. Thus, $(0, -\frac{1}{2})$ is the y -intercept. To find an x intercept, set $y = 0$. This gives $0 = \frac{1}{x-2} \Rightarrow 0(x-2) = \frac{1}{x-2}(x-2) \Rightarrow 0 = 1$, which has no solution. Thus, there are no x -intercepts.

Now for some ordered pairs that satisfy the formula $y = \frac{1}{x-2}$:

If we plot these points and connect them with a smooth curve, we would get the following graph:

x	y
-3	$-\frac{1}{5}$
-2	$-\frac{1}{4}$
-1	$-\frac{1}{3}$
0	$-\frac{1}{2}$
1	-1
2	Und.
3	1
4	$\frac{1}{2}$
5	$\frac{1}{3}$
6	$\frac{1}{4}$
7	$\frac{1}{5}$




What some students do at this point is to simply connect the points $(3, 1)$ and $(1, -1)$ with a straight line. Talk about jumping to conclusions!

Our domain of $\mathbb{R} - \{2\}$ implies that x cannot be 2 in this function; the straight line trick won't work.

So we agree that a major chunk of the graph is missing. How do we get a better picture of the graph? We try some x -values that are near 2:

$$\left(1\frac{1}{2}, -2\right) \quad \left(1\frac{3}{4}, -4\right) \quad \left(1\frac{7}{8}, -8\right) \quad \left(2\frac{1}{2}, 2\right) \quad \left(2\frac{1}{4}, 4\right) \quad \left(2\frac{1}{8}, 8\right)$$

Adding these points to our previous attempt at a graph gives us a much better picture:



This graph has some real cool **limits**. Suppose we let x approach ∞ . The y -values are positive (the curve is above the x -axis), but are getting smaller and smaller, approaching zero. Thus, as $x \rightarrow \infty$, $y \rightarrow 0$. Now let x approach $-\infty$. The y -values are negative but are rising toward zero. Therefore, as $x \rightarrow -\infty$, $y \rightarrow 0$.

The number 2 seems to be an interesting x -value. Although x can never be 2 in this function, it looks like the curve is getting closer and closer to the vertical line $x = 2$. In fact, if we let x approach 2 from the right, the curve is growing taller and taller, and so we have the limit: As $x \rightarrow 2^+$, $y \rightarrow \infty$. Now let x approach 2 from the left. This time the curve is dropping rapidly, toward negative infinity. This observation yields the limit: As $x \rightarrow 2^-$, $y \rightarrow -\infty$.

Let's summarize the four limits we've deduced:

$$\begin{array}{ll} \text{As } x \rightarrow \infty, y \rightarrow 0 & \text{As } x \rightarrow -\infty, y \rightarrow 0 \\ \text{As } x \rightarrow 2^+, y \rightarrow \infty & \text{As } x \rightarrow 2^-, y \rightarrow -\infty \end{array}$$

From these four limits we can conclude that the **range** of the function is all real numbers except 0. That is, the range is $\mathbb{R} - \{0\}$.

6

Do you see that as you move far to the right or far to the left, the curve gets closer and closer to the x -axis? We say that the line $y = 0$ (which is the x -axis) is a **horizontal asymptote**.

Now look at the region of the graph near $x = 2$. The curve gets closer and closer to the vertical line $x = 2$ (in fact, on both sides of the vertical line). We call the line $x = 2$ a **vertical asymptote**.

EXAMPLE 2: **Graph:** $y = \frac{2x-1}{x+2}$

Solution: First we find the **domain**. Recall that this function will be undefined when the denominator is zero, which occurs when $x = -2$. Thus, the domain is $\mathbb{R} - \{-2\}$.

Now let's explore the **intercepts**:

If $x = 0$, then $y = \frac{2(0)-1}{0+2} = -\frac{1}{2}$. There's a y -intercept at $(0, -\frac{1}{2})$.

If $y = 0$, then $0 = \frac{2x-1}{x+2} \Rightarrow 2x-1 = 0 \Rightarrow x = \frac{1}{2}$. So $(\frac{1}{2}, 0)$ is an x -intercept.

It's time for some more ordered pairs for this function. Use your calculator to verify each of the following:

$$(-1, -3) \quad (1, 0.33) \quad (3, 1) \quad (5, 1.29) \quad (10, 1.58)$$

$$(15, 1.71) \quad (20, 1.77) \quad (100, 1.95) \quad (1000, 1.995)$$

What's happening as x grows very large? It appears that y is approaching 2. That is, as $x \rightarrow \infty$, $y \rightarrow 2$.

Now we'll let x go the other direction:

$$(-3, 7) \quad (-5, 3.67) \quad (-10, 2.63) \quad (-20, 2.28)$$

$$(-100, 2.05) \quad (-1000, 2.01)$$


These points show that as $x \rightarrow -\infty$, $y \rightarrow 2$.

Finally, here are some ordered pairs for x 's near -2 (the only real number not in the domain):

$(-1.5, -8)$ $(-1.9, -48)$ $(-1.99, -498)$ Thus, as $x \rightarrow -2^+$, $y \rightarrow -\infty$.

$(-2.5, 12)$ $(-2.1, 52)$ $(-2.01, 502)$ Therefore, as $x \rightarrow -2^-$, $y \rightarrow \infty$.

Plotting as many of the calculated points as possible, and taking into account the four limits we've found, the following graph emerges:



We can determine the **range** of the function by looking at the graph and seeing that y can be any value except 2. So the range is $\mathbb{R} - \{2\}$.

We can now be reasonably sure of the **asymptotes**. Either by recalling the limits described above or by looking at the graph, we conclude that there's a vertical asymptote at $x = -2$ and a horizontal asymptote at $y = 2$.

EXAMPLE 3: **Graph:** $y = \frac{4}{1+x^2}$

Solution: Why is this function rational? Because it's the ratio P/Q of two polynomial functions: the constant polynomial $P(x) = 4$ and the quadratic polynomial $Q(x) = 1 + x^2$.

To find the **domain**, set the denominator to zero to see what's not in the domain: $1 + x^2 = 0$. This equation has no solution in \mathbb{R} , since solving it leads to $x = \pm\sqrt{-1}$, which are not real numbers. In fact, for any value of x , the quantity $1 + x^2$ is at least 1 (why?), so it certainly can't be zero. Since the denominator can never be zero, there's nothing to be excluded from the domain, and therefore the domain is \mathbb{R} . We can also figure that the graph will not have a **vertical asymptote**.

Let's check for y -axis **symmetry**: Replace x with $-x$:

$$y = \frac{4}{1+(-x)^2} = \frac{4}{1+x^2}$$

It's the same equation; our graph has y -axis symmetry. This means that we need only use x 's that are ≥ 0 -- that is, only points to the right of the y -axis need to be plotted. The graph on the left side of the y -axis will be a mirror image of the right side.

Now we seek the **intercepts**. Set $x = 0$ to get $y = 4$, and so the y -intercept is $(0, 4)$. Now set $y = 0$, giving

$$0 = \frac{4}{1+x^2} \Rightarrow 0(1+x^2) = \frac{4}{1+x^2}(1+x^2) \Rightarrow 0 = 4.$$

This absurd result indicates that the equation has no solution; hence, there are no x -intercepts.

It's time for some ordered pairs for this function:

$$(1, 2) \quad (2, 0.8) \quad (3, 0.4) \quad (4, 0.24) \quad (10, 0.04) \quad (200, 0.0001)$$


These points suggest the limit: As $x \rightarrow \infty, y \rightarrow 0$. This implies that $y = 0$ is a horizontal **asymptote**.

Here are some more ordered pairs, designed to see what happens as we approach the y -axis from the right:

$$(0.75, 2.56) \quad (0.5, 3.2) \quad (0.25, 3.76) \quad (0.1, 3.96) \quad (0.02, 3.998)$$

These points give us the limit: As $x \rightarrow 0^+, y \rightarrow 4$.

If we plot all the points calculated so far, and if we recall the y -axis symmetry, we get the following graph:



We determined at the outset that the domain of this rational function is \mathbb{R} . Is it clear from the graph that this is indeed the case?


The graph shows us that the **range** of the function is all real numbers between 0 and 4, excluding the 0 but including the 4. This set can be written $\{y \in \mathbb{R} \mid 0 < y \leq 4\}$, or in interval notation as $(0, 4]$.

Homework

5. Consider the rational function in Example 2. Without referring to the graph, prove that y can have the value 2.01, but y can never have the value 2. Also without referring to the graph, verify that the graph does not have y -axis symmetry.
6. Find the domain:
- a. $y = \frac{2x+7}{9}$ b. $f(x) = \frac{2x-3}{4+x}$
- c. $g(x) = \frac{3x}{2x-10}$ d. $R(x) = \frac{x-1}{-7x+4}$
- e. $f(x) = \frac{x^2-9}{x^2-100}$ f. $g(x) = \frac{8x-16}{x^2+25}$
7. Find the intercepts:
- a. $f(x) = \frac{x-4}{x-2}$ b. $y = \frac{3}{5x-15}$
- c. $y = \frac{2x+1}{x-3}$ d. $g(x) = \frac{5-x}{6x+1}$
8. Find the asymptotes:
- a. $R(x) = \frac{8x+1}{4x-4}$ b. $y = \frac{2x-3}{2x+1}$
- c. $y = \frac{3x-7}{x+2}$ d. $h(x) = \frac{2x+7}{4x-4}$
9. Find the domain, the intercepts, and the asymptotes of

$$y = \frac{1}{4+x^2}.$$
10. Perform a complete analysis of the function $y = \frac{2}{x-3}$.
11. Perform a complete analysis of the function $y = \frac{3x-5}{x-2}$.

12. Perform a complete analysis of the function $y = \frac{2}{2+x^2}$.
13. Perform a complete analysis of the function $y = \frac{-1}{x+1}$.
14. Consider the graph



Explain why the horizontal line $y = 0$ (that is, the x -axis) is a horizontal asymptote for the curve.

Practice Problems

15. Explain why $f(x) = \sqrt{7}x^{10} + \pi x^7 - 6x - 1$ is a polynomial. What is its degree?
16. Explain why $y = 3x^5 - \sqrt{x} + \pi$ is not a polynomial.
17. a. A horizontal line (is, is not) a polynomial.
 b. The function $y = \frac{1}{x}$ (is, is not) a polynomial.
 c. What is the degree of the polynomial $y = 7x - \pi$?
 d. Is a circle a polynomial?

18. Consider the rational function $y = \frac{7}{2x-8}$.
- Find the domain.
 - Find all the intercepts.
 - Find all the asymptotes.
 - Calculate y if $x = 4.1$.
19. Find all the intercepts and asymptotes of $r(x) = \frac{8x+6}{2x-3}$, and graph.
20. Graph $y = \frac{-2x-2}{x-1}$. As $x \rightarrow 1^+$, $y \rightarrow \underline{\hspace{1cm}}$. As $x \rightarrow 1^-$, $y \rightarrow \underline{\hspace{1cm}}$.
As $x \rightarrow \infty$, $y \rightarrow \underline{\hspace{1cm}}$. As $x \rightarrow -\infty$, $y \rightarrow \underline{\hspace{1cm}}$.
21. Graph $y = \frac{5}{2+x^2}$. Discuss domain, range, symmetry, and asymptotes. As $x \rightarrow \infty$, $y \rightarrow \underline{\hspace{1cm}}$. As $x \rightarrow -\infty$, $y \rightarrow \underline{\hspace{1cm}}$.
As $x \rightarrow 0^+$, $y \rightarrow \underline{\hspace{1cm}}$. As $x \rightarrow 0^-$, $y \rightarrow \underline{\hspace{1cm}}$.
22. True/False:
- $y = \sqrt[3]{7}x^{10} - \pi x^3 + \sqrt{2}$ is a polynomial.
 - $y = \frac{1}{x^5}$ is a polynomial.
 - The graph of $f(x) = \frac{1}{2x+10}$ has a vertical asymptote at $x = -5$.
 - The graph of $g(x) = \frac{10x+9}{5x-11}$ has a horizontal asymptote at $y = 10$.
 - The range of the function $y = \frac{6}{1+x^2}$ is $(0, 6]$.
 - The above function has x -axis symmetry.
 - The above function has a domain of $\mathbb{R} - \{\pm 1\}$.
 - For the graph of $y = \frac{3x+1}{x-\pi}$, as $x \rightarrow \infty$, $y \rightarrow 3$.

Solutions

1. All coefficients are from \mathbb{R} , and all exponents are from \mathbb{W} .
2. The middle term is $\sqrt{2}x^{1/2}$, and $\frac{1}{2} \notin \mathbb{W}$.
3. a. 0 b. 3 c. 10 d. 1
4. a. \mathbb{R} b. False c. True d. False
5. $y = \frac{2x-1}{x+2} \Rightarrow 2.01 = \frac{2x-1}{x+2} \Rightarrow 2.01x + 4.02 = 2x - 1 \Rightarrow x = -502$.
So, $(-502, 2.01)$ is on the graph, and indeed y can be 2.01.

Now let's pretend that y could be 2; then

$2 = \frac{2x-1}{x+2} \Rightarrow 2x + 4 = 2x - 1 \Rightarrow 4 = -1 \Rightarrow$ no solution. Thus, there is no x which will make $y = 2$.

To test for y -axis symmetry, replace x with $-x$:

$y = \frac{2(-x)-1}{-x+2} \Rightarrow y = \frac{-2x-1}{-x+2}$, which is not the original formula, nor can it be made to be the original formula. You test for origin symmetry.


6. a. \mathbb{R} b. $\mathbb{R} - \{-4\}$ c. $\mathbb{R} - \{5\}$ d. $\mathbb{R} - \left\{\frac{4}{7}\right\}$ e. $\mathbb{R} - \{\pm 10\}$ f. \mathbb{R}
7. a. $(4, 0)$ $(0, 2)$ b. $(0, -\frac{1}{5})$ c. $(-\frac{1}{2}, 0)$ $(0, -\frac{1}{3})$ d. $(5, 0)$ $(0, 5)$
8. a. $x = 1$ $y = 2$ b. $x = -\frac{1}{2}$ $y = 1$ c. $x = -2$ $y = 3$ d. $x = 1$ $y = \frac{1}{2}$

9. Since the only way the formula can be messed up is by dividing by 0, and since the denominator can never be zero (verify this yourself), the domain is \mathbb{R} .

Setting $x = 0$ gives a y -value of $1/4$, so the y -intercept is $(0, \frac{1}{4})$. If you set $y = 0$, you'll get no solution for y . Thus, there is no x -intercept.

There are no vertical asymptotes, since the denominator is never zero. Letting x approach either ∞ or $-\infty$, y approaches 0. Thus, a horizontal asymptote is $y = 0$ (the x -axis).

10.



Domain = $\mathbb{R} - \{3\}$

x -int: none

y -int: $(0, -\frac{2}{3})$

As $x \rightarrow 3^+$, $y \rightarrow \infty$

As $x \rightarrow 3^-$, $y \rightarrow -\infty$

As $x \rightarrow \infty$, $y \rightarrow 0$

As $x \rightarrow -\infty$, $y \rightarrow 0$


vert asy: $x = 3$

horiz asy: $y = 0$

Range = $\mathbb{R} - \{0\}$

No symmetry

11.



Domain = $\mathbb{R} - \{2\}$

x -int: $(\frac{5}{3}, 0)$

y -int: $(0, \frac{5}{2})$

As $x \rightarrow 2^+$, $y \rightarrow \infty$

As $x \rightarrow 2^-$, $y \rightarrow -\infty$

As $x \rightarrow \infty$, $y \rightarrow 3$

As $x \rightarrow -\infty$, $y \rightarrow 3$


vert asy: $x = 2$

horiz asy: $y = 3$

Range = $\mathbb{R} - \{3\}$


No symmetry

12.



Domain = \mathbb{R} x -int: none y -int: (0, 1) Symmetry: y -axis
 As $x \rightarrow \infty, y \rightarrow 0$ As $x \rightarrow -\infty, y \rightarrow 0$
 vert asy: none horiz asy: $y = 0$
 maximum point at (0, 1)
 Range = $\{y \in \mathbb{R} \mid 0 < y \leq 1\} = (0, 1]$

13.




Domain = $\mathbb{R} - \{-1\}$
 x -int: none
 y -int: (0, -1)
 As $x \rightarrow -1^+, y \rightarrow -\infty$
 As $x \rightarrow -1^-, y \rightarrow \infty$
 As $x \rightarrow \infty, y \rightarrow 0$
 As $x \rightarrow -\infty, y \rightarrow 0$
 vert asy: $x = -1$
 horiz asy: $y = 0$
 Range = $\mathbb{R} - \{0\}$
 No symmetry


14. Because of the limit: As $x \rightarrow \infty, y \rightarrow 0$. Even though the graph intersects its own horizontal asymptote infinitely often, the curve nevertheless continues to get closer and closer to the x -axis (the line $y = 0$), and this is ultimately what is meant by a horizontal asymptote.

15. f is a polynomial because the coefficients are real numbers and the exponents (the 10, 7 and 1) are whole numbers. Its degree is 10.


16. Look at the middle term; it can be written as $x^{1/2}$, a term whose exponent is not from the whole numbers.
17. a. is b. is not ($1/x = x^{-1}$) c. 1 d. It's not even a function, let alone the special function called a polynomial.
18. a. $\mathbb{R} - \{4\}$ b. $(0, -7/8)$ c. $x = 4$ and $y = 0$ d. 35
19. Intercepts: $(0, -2)$ and $(-\frac{3}{4}, 0)$; vert asy: $x = \frac{3}{2}$; horiz asy: $y = 4$



20.

Limits: $-\infty$; ∞ ; -2 ; -2

21.


Domain = \mathbb{R} Range = $(0, \frac{5}{2}]$ y -axis symmetry

No vert asy

Horiz asy: $y = 0$ Limits: $0; 0; \frac{5}{2}; \frac{5}{2}$

22. a. T b. F c. T d. F e. T f. F g. F h. T

“ The universe cannot be read until we have learned the language and become familiar with the characters in which it is written. It is written in mathematical language.”



Galileo Galiliei